#### **Patient Details**

## Constipation & Constipation Predominant IBS (IBS-C)

(Abdominal Pain / Bloating / Constipation)

The majority of patients meeting the diagnostic criteria below, without 'red flag indicators' and with normal blood tests do not require referral to secondary care.

1) Linaclotide 290mcg od (to soft stool + for pain) or

bloating and to increase the defaecation frequency)

Review in 1 Month to check for initial response (Full benefit may

2) Prucalopride 2mg od (NICE Approved for pain,

not be reached for up to 3m)

#### Typical functional symptoms

- Symptoms present for at least 3-6 months
- Abdominal pain or discomfort relieved by defecation OR
- Associated with altered bowel frequency or form
- <3 bowel movements per week + one or more of the following:
- Altered stool passage (straining, urgency, incomplete evacuation)
- Self medications / Prescribed on either a Laxative or Antispasmodic
- Abdominal bloating (distension, tension or hardness
- · Symptoms made worse by eating
- Passage of mucus
- Lethargy

and review in 6

months

### **Red Flags Symptoms:**

- Unintentional & unexplained weight loss
- Rectal bleeding
- Anaemia

oor Response

- Abdominal or rectal masses
- Change in bowel habit looser / frequent stools
- >60y, persisting for >6w with loose more frequent stools
- Unexplained raised ESR or CRP
- FH of bowel / ovarian cancer (consider pelvic US+Ca125)
- +ve Faecal Calprotectin (>100)

## 

# Secondary Care Referral

Patients with <u>RED</u> flag indicators <u>MUST</u> be referred to secondary care for assessment +/- endoscopic review

Tests expected prior to referral Please list results below

- FBC
- Ferritin
- Coeliac screen
- TFTs
- Faecal calprotectin
- +/- AXR